


*Physics perspectives with heavy ions at the High Luminosity
- LHC and beyond*

Stefan Floerchinger (Heidelberg U.)

LHCP conference, Bologna, 08.06.2018



UNIVERSITÄT
HEIDELBERG
ZUKUNFT
SEIT 1386




Heavy ions at the HL-LHC

Ongoing discussion, see for example:

- Jan-Fiete Grosse-Oetringhaus, talk at *Workshop on the physics of HL-LHC*, 30.10.2017: <https://indico.cern.ch/event/647676/timetable/>
- Andrea Dainese, talk at *ECFA High Luminosity LHC Experiments Workshop*, 04.10.2016: <https://indico.cern.ch/event/524795/timetable/>
- J. M. Jowett, M. Schaumann and R. Versteegen, *Heavy-Ion Operation of HL-LHC*: <https://cds.cern.ch/record/1977371>
- Antonio Uras, *Heavy-Ions at the High-Luminosity LHC*: <http://inspirehep.net/record/1589642>
- preparation of a CERN yellow report chapter on *Heavy ions at the HL-LHC*, working group meeting: <https://indico.cern.ch/event/717641/>
- existing CERN yellow report chapter on *Heavy Ions at the Future Circular Collider*: <http://inspirehep.net/record/1455787?ln=de>

I will not attempt to reflect the full ongoing discussion, but rather present my own point of view (as a theorist).


Little bangs in the laboratory



A great challenge

- quantum fields at finite energy density and temperature
- fundamental gauge theory: QCD
- strongly interacting
- non-equilibrium dynamics
- experimentally driven field of research
- big motivation for theory development

Fluid dynamics



- long distances, long times or strong enough interactions
- matter or quantum fields form a fluid!
- needs **macroscopic** fluid properties
 - thermodynamic equation of state $p(T, \mu)$
 - shear viscosity $\eta(T, \mu)$
 - bulk viscosity $\zeta(T, \mu)$
 - heat conductivity $\kappa(T, \mu)$
 - relaxation times, ...
- *ab initio* calculation of fluid properties difficult but fixed by **microscopic** properties in \mathcal{L}_{QCD}

Relativistic fluid dynamics

Energy-momentum tensor and conserved current

$$T^{\mu\nu} = \epsilon u^\mu u^\nu + (p + \pi_{\text{bulk}}) \Delta^{\mu\nu} + \pi^{\mu\nu}$$

$$N^\mu = n u^\mu + \nu^\mu$$

- tensor decomposition using fluid velocity u^μ , $\Delta^{\mu\nu} = g^{\mu\nu} + u^\mu u^\nu$
- thermodynamic equation of state $p = p(T, \mu)$

Covariant **conservation laws** $\nabla_\mu T^{\mu\nu} = 0$ and $\nabla_\mu N^\mu = 0$ imply

- equation for **energy density** ϵ

$$u^\mu \partial_\mu \epsilon + (\epsilon + p + \pi_{\text{bulk}}) \nabla_\mu u^\mu + \pi^{\mu\nu} \nabla_\mu u_\nu = 0$$

- equation for **fluid velocity** u^μ

$$(\epsilon + p + \pi_{\text{bulk}}) u^\mu \nabla_\mu u^\nu + \Delta^{\nu\mu} \partial_\mu (p + \pi_{\text{bulk}}) + \Delta^\nu{}_\alpha \nabla_\mu \pi^{\mu\alpha} = 0$$

- equation for **particle number density** n

$$u^\mu \partial_\mu n + n \nabla_\mu u^\mu + \nabla_\mu \nu^\mu = 0$$

Constitutive relations

Second order relativistic fluid dynamics:

- equation for **shear stress** $\pi^{\mu\nu}$

$$\tau_{\text{shear}} P^{\rho\sigma}_{\alpha\beta} u^\mu \nabla_\mu \pi^{\alpha\beta} + \pi^{\rho\sigma} + 2\eta P^{\rho\sigma\alpha}_{\beta} \nabla_\alpha u^\beta + \dots = 0$$

with **shear viscosity** $\eta(T, \mu)$

- equation for **bulk viscous pressure** π_{bulk}

$$\tau_{\text{bulk}} u^\mu \partial_\mu \pi_{\text{bulk}} + \pi_{\text{bulk}} + \zeta \nabla_\mu u^\mu + \dots = 0$$



with **bulk viscosity** $\zeta(T, \mu)$

- equation for **baryon diffusion current** ν^μ

$$\tau_{\text{heat}} \Delta^\alpha_\beta u^\mu \nabla_\mu \nu^\beta + \nu^\alpha + \kappa \left[\frac{nT}{\epsilon + p} \right]^2 \Delta^{\alpha\beta} \partial_\beta \left(\frac{\mu}{T} \right) + \dots = 0$$

with **heat conductivity** $\kappa(T, \mu)$

Thermodynamics of QCD



[Borsányi *et al.* (2016)], similar Bazavov *et al.* (2014)

[Bazavov *et al.* (2017)], similar Bellwied *et al.* (2015)]

- thermodynamic equation of state $p(T)$ rather well understood now
- also moments of conserved charges like

$$\chi_2^B = \frac{\langle (N_B - N_{\bar{B}})^2 \rangle}{VT^3}$$


and higher order understood

- progress in computing power


Quantum fields and information

- surprising relations between quantum field theory and information theory
- well understood in thermal equilibrium
- currently investigated out-of-equilibrium
- fluid dynamics / entanglement entropy / black hole physics (AdS/CFT)
- shear viscosity to entropy density ratio $\eta/s \geq \hbar/(4\pi k_B)$

[Kovtun, Son, Starinets (2003)]




[Ryu, Takayanagi (2006)]



[Berges, Floerchinger, Venugopalan (2017)]

Non-central collisions



- pressure gradients larger in reaction plane
- leads to larger fluid velocity in this direction
- more particles fly in this direction
- can be quantified in terms of elliptic flow v_2
- particle distribution

$$\frac{dN}{d\phi} = \frac{N}{2\pi} \left[1 + 2 \sum_m v_m \cos(m(\phi - \psi_R)) \right]$$


- symmetry $\phi \rightarrow \phi + \pi$ implies $v_1 = v_3 = v_5 = \dots = 0$.

Two-particle correlation function

- normalized two-particle correlation function

$$C(\phi_1, \phi_2) = \frac{\langle \frac{dN}{d\phi_1} \frac{dN}{d\phi_2} \rangle_{\text{events}}}{\langle \frac{dN}{d\phi_1} \rangle_{\text{events}} \langle \frac{dN}{d\phi_2} \rangle_{\text{events}}} = 1 + 2 \sum_m v_m^2 \cos(m(\phi_1 - \phi_2))$$


- surprisingly v_2, v_3, v_4, v_5 and v_6 are all non-zero!





[ALICE 2011, similar results from CMS, ATLAS, Phenix, Star]

Event-by-event fluctuations

- deviations from symmetric initial energy density distribution from event-by-event fluctuations
- one example is Glauber model




Big bang – little bang analogy



- cosmol. scale: $\text{Mpc} = 3.1 \times 10^{22} \text{ m}$
- Gravity + QED + Dark sector
- one big event
- initial conditions not directly accessible
- all information must be reconstructed from final state
- dynamical description as a fluid
- fluctuating initial state
- nuclear scale: $\text{fm} = 10^{-15} \text{ m}$
- QCD
- very many events

Similarities to cosmological fluctuation analysis



- fluctuation spectrum contains info from early times
- detailed correlation functions are compared to theory
- can lead to detailed understanding of evolution
- Mode-by-mode fluid dynamics for heavy ion collisions

[Floerchinger, Wiedemann (2014)]

The dark matter fluid

- **high energy nuclear collisions**

$$\mathcal{L}_{\text{QCD}} \rightarrow \text{fluid properties}$$

- **late time cosmology**

$$\text{fluid properties} \rightarrow \mathcal{L}_{\text{dark matter}}$$

- until direct detection of dark matter it can only be observed via gravity


$$G^{\mu\nu} = 8\pi G_{\text{N}} T^{\mu\nu}$$

so all we can access is

$$T_{\text{dark matter}}^{\mu\nu}$$

- strong motivation to study heavy ion collisions and cosmology together!


Collective behavior in large and small systems



- flow coefficients from higher order cumulants $v_2\{n\}$ agree:
→ collective behavior
- elliptic flow signals also in **pPb** and **pp** !
- can fluid approximation work for pp collisions?

Questions and puzzles



- how universal are collective flow and fluid dynamics?
 - as a limit of kinetic theory / perturbation theory / multi-parton interactions
 - non-perturbative understanding / entanglement
- what determines density distribution of a proton?
 - constituent quarks or interacting gluon cloud?
 - generalized PDFs
- more elementary collision systems? **[News at Quark Matter 2018!]**



- role of electromagnetic fields and vorticity for fluid dynamics
- role of quantum anomalies (e. g. chiral magnetic effect)



Chemical freeze-out

[Andronic, Braun-Munzinger, Redlich, Stachel (2017)]




- chemical freeze-out close to chiral crossover transition for large \sqrt{s}
- chiral transition should be visible in higher moments $\langle (N_B - N_{\bar{B}})^n \rangle$
- traces of the evolving chiral condensate / pion condensate ?
- more insights at large μ_B expected from FAIR

Quarkonium and how it gets modified



- all Υ states are suppressed by medium effects, excited states even more
- more detailed understanding of heavy quark bound states in a medium
- also at LHC: regeneration and flow of charmed mesons
- future: also bottom

Jet quenching





- asymmetry between reconstructed jet energies

$$A_J = \frac{E_{T1} - E_{T2}}{E_{T1} + E_{T2}}, \quad \Delta\phi > \pi/2$$

- partons/jets lose energy to the quark gluon plasma
- jet structure can be investigated in detail
- more possible: b -jets, t -jets
- interplay of microscopic partons / jets and macroscopic QCD fluid

Light-by-light scattering

[ATLAS, Nature Phys. 13, 852 (2017)]



- ultra-peripheral ion collisions produce strong electromagnetic fields
- beam of quasi-real photons (equivalent photon approximation)
- Halpern scattering $\gamma\gamma \rightarrow \gamma\gamma$ observed, more detailed studies possible
- also ultra-peripheral: nuclear PDFs

Theory development


- many interesting experimental results available or in reach
- precise studies need interplay of theory and experiment
- **more dedicated theory development needed**
- **we need to develop and maintain a standard model**
- heavy ion collisions and QCD dynamics can be understood much better !

Plans for heavy ions at runs 2-4 at the LHC

[J.-F. Grosse-Oetringhaus, CERN, 30.10.2017]

- Run 2:
 - Pb-Pb: few nb^{-1} (0.7 nb^{-1} in 2015, $\sim 1 \text{ nb}^{-1}$ in 2018) at $\sqrt{s_{\text{NN}}} = 5 \text{ TeV}$
 - p-Pb at 5 and 8 TeV (185 nb^{-1} in 2016)
 - pp reference at Pb-Pb energy (5 TeV, Nov 2017)
- LS2:
 - LHC injector upgrades; bunch spacing reduced to 50 ns
 - Pb-Pb interaction rate up to 50 kHz (now <10 kHz)
 - Experiments' upgrades (also LS3)
- Runs 3+4:
 - Request for **Pb-Pb: $>10 \text{ nb}^{-1}$**
(ALICE: 10 nb^{-1} at 0.5T + 3 nb^{-1} at 0.2T)
 - In line with projections by machine:
3.1 $\text{nb}^{-1}/\text{month}$ (Chamonix 2017)

$$\sigma_{\text{hadr,PbPb}} = 8 \text{ barn !}$$







HL-LHC for heavy ions begins in Run 3 !



Detector Upgrades


most relevant to heavy-ion physics

- **ALICE** (LS2)
 - New inner tracker: **precision and efficiency at low p_T**
 - New pixel forward muon tracker: **precise tracking and vertexing for μ**
 - TPC upgrade + readout + online data reduction **x100 faster readout (continuous)**
- **ATLAS** (LS2/LS3)
 - Fast tracking trigger (LS2): **high-multiplicity tracking**
 - Calorimeter and muon upgrades (LS2): **electron, γ , muon triggers**
 - ZDC replacement planned (LS2): **radiation hardness, granularity**
 - Completely new tracker (LS3): **tracking and b-tag up to $\eta=4$**
- **CMS** (mainly LS3)
 - Extension of forward muon system (LS2): **muon acceptance**
 - Completely new tracker (LS3): **tracking and b-tag up to $\eta=4$**
 - Upgrade forward calorimeter (LS3): **forward jets in HL**
- **LHCb** (LS2)
 - Triggerless readout, full software trigger, higher granularity detectors: **impact on tracking performance in Pb-Pb being studied**
 - Fixed-target programme with SMOG + possible extensions



Higher energies

[Dainese, Wiedemann (ed.) et al. (2017)]



Quantity	Pb–Pb 2.76 TeV	Pb–Pb 5.5 TeV	Pb–Pb 39 TeV
$dN_{\text{ch}}/d\eta$ at $\eta = 0$	1600	2000	3600
Total N_{ch}	17000	23000	50000
$dE_T/d\eta$ at $\eta = 0$	1.8–2.0 TeV	2.3–2.6 TeV	5.2–5.8 TeV
Homogeneity volume	5000 fm ³	6200 fm ³	11000 fm ³
Decoupling time	10 fm/c	11 fm/c	13 fm/c
ε at $\tau = 1$ fm/c	12–13 GeV/fm ³	16–17 GeV/fm ³	35–40 GeV/fm ³

Larger collision energy

- higher initial energy density and temperature
- higher multiplicity N_{ch}
- larger lifetime and volume of fireball
- better probes of collective physics
- thermal charm quarks
- more hard probes

A dedicated detector for low p_T ?

- advances in detector technology might allow to construct dedicated detector for low p_T spectrum
- down to $p_T \approx 10 \text{ MeV} \approx \frac{1}{20 \text{ fm}}$?
- low momentum di-leptons
 - excellent understanding of charmonia and bottomonia (P-wave)
- probe macroscopic properties of QCD fluid: very soft pions, kaons, protons, di-leptons
 - dynamics of chiral symmetry restoration
 - pion condensates / disoriented chiral condensates ?
- understand thermalization and dissipation in detail
 - spectrum also at $p_T \ll T_{\text{kinetic freeze-out}} \approx 120 \text{ MeV}$

Conclusions

- high energy nuclear collisions produce a relativistic QCD fluid!
- interesting parallels between cosmology and heavy ion collisions
- heavy ion collisions provide chance to understand a relativistic fluid from first principles
- experimental hints for collective flow also in pPb and pp collisions
- QCD fluid can be understood in much more detail with combined effort of theory and experiment!
- *I had to skip many interesting topics, please see also other presentations mentioned on the first slide.*