


Particle production from expanding QCD strings and its quantum simulation

Stefan Floerchinger (Heidelberg U.)

Quantum Systems in Extreme Conditions
Heidelberg University, 25.09.2019




UNIVERSITÄT
HEIDELBERG
ZUKUNFT
SEIT 1386



The thermal model puzzle


- elementary particle collision experiments such as $e^+ e^-$ collisions show some thermal-like features
- particle multiplicities well described by thermal model



[Becattini, Casterina, Milov & Satz, EPJC 66, 377 (2010)]

- conventional thermalization by collisions unlikely
- more thermal-like features difficult to understand in PYTHIA
[Fischer, Sjöstrand (2017)]
- alternative explanations needed

QCD strings




- particle production from QCD strings
- Lund string model (e. g. PYTHIA)
- different regions in a string are entangled
- subinterval A is described by reduced density matrix

$$\rho_A = \text{Tr}_B \rho$$

- reduced density matrix is of mixed state form
- could this lead to thermal-like effects?

Entropy and entanglement

- consider a split of a quantum system into two $A + B$



- reduced density operator for system A

$$\rho_A = \text{Tr}_B\{\rho\}$$

- entropy associated with subsystem A : **entanglement entropy**

$$S_A = -\text{Tr}_A\{\rho_A \ln \rho_A\}$$

- globally pure** state $S = 0$ can be **locally mixed** $S_A > 0$
- coherent information** $I_{B \rangle A} = S_A - S$ can be **positive**

Microscopic model

- QCD in 1+1 dimensions described by 't Hooft model

$$\mathcal{L} = -\bar{\psi}_i \gamma^\mu (\partial_\mu - ig \mathbf{A}_\mu) \psi_i - m_i \bar{\psi}_i \psi_i - \frac{1}{2} \text{tr} \mathbf{F}_{\mu\nu} \mathbf{F}^{\mu\nu}$$

- fermionic fields ψ_i with sums over flavor species $i = 1, \dots, N_f$
- $\text{SU}(N_c)$ gauge fields $\mathbf{A}_\mu$ with field strength tensor $\mathbf{F}_{\mu\nu}$
- gluons are not dynamical in two dimensions
- gauge coupling g has dimension of mass
- non-trivial, interacting theory, cannot be solved exactly
- spectrum of excitations known for $N_c \rightarrow \infty$ with $g^2 N_c$ fixed
['t Hooft (1974)]

Schwinger model

- QED in 1+1 dimension

$$\mathcal{L} = -\bar{\psi}_i \gamma^\mu (\partial_\mu - iqA_\mu) \psi_i - m_i \bar{\psi}_i \psi_i - \frac{1}{4} F_{\mu\nu} F^{\mu\nu}$$

- geometric confinement
- U(1) charge related to string tension $q = \sqrt{2\sigma}$
- for single fermion one can **bosonize theory** exactly
[Coleman, Jackiw, Susskind (1975)]

$$S = \int d^2x \sqrt{g} \left\{ -\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \frac{1}{2} M^2 \phi^2 - \frac{m q e^\gamma}{2\pi^{3/2}} \cos(2\sqrt{\pi}\phi + \theta) \right\}$$

- Schwinger bosons are dipoles $\phi \sim \bar{\psi}\psi$
- scalar mass related to U(1) charge by $M = q/\sqrt{\pi} = \sqrt{2\sigma/\pi}$
- massless Schwinger model $m = 0$ leads to free bosonic theory


Transverse coordinates

- so far dynamics strictly confined to 1+1 dimensions
- transverse coordinates may fluctuate, can be described by Nambu-Goto action ($h_{\mu\nu} = \partial_\mu X^m \partial_\nu X_m$)

$$\begin{aligned} S_{\text{NG}} &= \int d^2x \sqrt{-\det h_{\mu\nu}} \{-\sigma + \dots\} \\ &\approx \int d^2x \sqrt{g} \left\{ -\sigma - \frac{\sigma}{2} g^{\mu\nu} \partial_\mu X^i \partial_\nu X^i + \dots \right\} \end{aligned}$$

- two additional, massless, bosonic degrees of freedom corresponding to transverse coordinates X^i with $i = 1, 2$

Expanding string solution 1



- external quark-anti-quark pair on trajectories $z = \pm t$
- coordinates: Bjorken time $\tau = \sqrt{t^2 - z^2}$, rapidity $\eta = \text{arctanh}(z/t)$
- metric $ds^2 = -d\tau^2 + \tau^2 d\eta^2$
- symmetry with respect to longitudinal boosts $\eta \rightarrow \eta + \Delta\eta$

Expanding string solution 2

- Schwinger boson field depends only on τ

$$\bar{\phi} = \bar{\phi}(\tau)$$

- equation of motion

$$\partial_\tau^2 \bar{\phi} + \frac{1}{\tau} \partial_\tau \bar{\phi} + M^2 \bar{\phi} = 0.$$

- Gauss law: electric field $E = q\phi/\sqrt{\pi}$ must approach the U(1) charge of the external quarks $E \rightarrow q_e$ for $\tau \rightarrow 0_+$

$$\bar{\phi}(\tau) \rightarrow \frac{\sqrt{\pi}q_e}{q} \quad (\tau \rightarrow 0_+)$$

- solution of equation of motion [Loshaj, Kharzeev (2011)]

$$\bar{\phi}(\tau) = \frac{\sqrt{\pi}q_e}{q} J_0(M\tau)$$

Gaussian states

- theories with quadratic action often have Gaussian density matrix
- fully characterized by field expectation values

$$\bar{\phi}(x) = \langle \phi(x) \rangle, \quad \bar{\pi}(x) = \langle \pi(x) \rangle$$

and connected two-point correlation functions, e. g.

$$\langle \phi(x)\phi(y) \rangle_c = \langle \phi(x)\phi(y) \rangle - \bar{\phi}(x)\bar{\phi}(y)$$

- if ρ is Gaussian, also reduced density matrix ρ_A is Gaussian

Entanglement entropy for Gaussian state

- entanglement entropy of Gaussian state in region A
[Berges, Floerchinger, Venugopalan, JHEP 1804 (2018) 145]


$$S_A = \frac{1}{2} \text{Tr}_A \{ D \ln(D^2) \}$$

- operator trace over region A only
- matrix of correlation functions

$$D(x, y) = \begin{pmatrix} -i\langle\phi(x)\pi(y)\rangle_c & i\langle\phi(x)\phi(y)\rangle_c \\ -i\langle\pi(x)\pi(y)\rangle_c & i\langle\pi(x)\phi(y)\rangle_c \end{pmatrix}$$

- involves connected correlation functions of field $\phi(x)$ and canonically conjugate momentum field $\pi(x)$
- expectation value $\bar{\phi}$ does not appear explicitly
- coherent states and vacuum have equal entanglement entropy S_A


Rapidity interval



- consider rapidity interval $(-\Delta\eta/2, \Delta\eta/2)$ at fixed Bjorken time τ
- entanglement entropy does not change by unitary time evolution with endpoints kept fixed
- can be evaluated equivalently in interval $\Delta z = 2\tau \sinh(\Delta\eta/2)$ at fixed time $t = \tau \cosh(\Delta\eta/2)$
- need to solve eigenvalue problem with correct **boundary conditions**

Bosonized massless Schwinger model

- entanglement entropy understood numerically for free massive scalars [Casini, Huerta (2009)]
- entanglement entropy density $dS/d\Delta\eta$ for bosonized massless Schwinger model ($M = \frac{q}{\sqrt{\pi}}$)



Conformal limit

- For $M\tau \rightarrow 0$ one has conformal field theory limit
[Holzhey, Larsen, Wilczek (1994)]

$$S(\Delta z) = \frac{c}{3} \ln (\Delta z/\epsilon) + \text{constant}$$

with small length ϵ acting as UV cutoff.

- Here this implies

$$S(\tau, \Delta\eta) = \frac{c}{3} \ln (2\tau \sinh(\Delta\eta/2)/\epsilon) + \text{constant}$$

- Conformal charge $c = 1$ for free massless scalars or Dirac fermions.
- Additive constant not universal but entropy density is

$$\begin{aligned} \frac{\partial}{\partial \Delta\eta} S(\tau, \Delta\eta) &= \frac{c}{6} \coth(\Delta\eta/2) \\ &\rightarrow \frac{c}{6} \quad (\Delta\eta \gg 1) \end{aligned}$$

- Entropy becomes extensive in $\Delta\eta$!

Universal entanglement entropy density

- for very early times “Hubble” expansion rate dominates over masses and interactions

$$H = \frac{1}{\tau} \gg M = \frac{q}{\sqrt{\pi}}, m$$

- theory dominated by free, massless fermions
- universal entanglement entropy density

$$\frac{dS}{d\Delta\eta} = \frac{c}{6}$$

with conformal charge c

- for QCD in 1+1 D (gluons not dynamical, no transverse excitations)

$$c = N_c \times N_f$$

- from fluctuating transverse coordinates (Nambu-Goto action)

$$c = N_c \times N_f + 2 \approx 9 + 2 = 11$$

Temperature and entanglement entropy

- for conformal fields, entanglement entropy has also been calculated at non-zero temperature.
- for static interval of length L [Korepin (2004); Calabrese, Cardy (2004)]

$$S(T, l) = \frac{c}{3} \ln \left(\frac{1}{\pi T \epsilon} \sinh(\pi L T) \right) + \text{const}$$


- compare this to our result in expanding geometry

$$S(\tau, \Delta\eta) = \frac{c}{3} \ln \left(\frac{2\tau}{\epsilon} \sinh(\Delta\eta/2) \right) + \text{const}$$

- expressions agree for $L = \tau\Delta\eta$ (with metric $ds^2 = -d\tau^2 + \tau^2 d\eta^2$) and time-dependent temperature

$$T = \frac{1}{2\pi\tau}$$

Modular or entanglement Hamiltonian 1



- conformal field theory
- hypersurface Σ with boundary on the intersection of two light cones
- reduced density matrix [Casini, Huerta, Myers (2011), Arias, Blanco, Casini, Huerta (2017), see also Candelas, Dowker (1979)]

$$\rho_A = \frac{1}{Z_A} e^{-K}, \quad Z_A = \text{Tr} e^{-K}$$

- modular or entanglement Hamiltonian K

Modular or entanglement Hamiltonian 2

- modular or entanglement Hamiltonian is **local expression**

$$K = \int_{\Sigma} d\Sigma_{\mu} \xi_{\nu}(x) T^{\mu\nu}(x).$$

- energy-momentum tensor $T^{\mu\nu}(x)$ of excitations
- vector field


$$\xi^{\mu}(x) = \frac{2\pi}{(q-p)^2} [(q-x)^{\mu}(x-p)(q-p) \\ + (x-p)^{\mu}(q-x)(q-p) - (q-p)^{\mu}(x-p)(q-x)]$$

end point of future light cone q , starting point of past light cone p

- inverse temperature and fluid velocity

$$\xi^{\mu}(x) = \beta^{\mu}(x) = \frac{u^{\mu}(x)}{T(x)}$$

Modular or entanglement Hamiltonian 3



- for $\Delta\eta \rightarrow \infty$: fluid velocity in τ -direction, τ -dependent temperature

$$T(\tau) = \frac{\hbar}{2\pi\tau}$$



- **Entanglement between different rapidity intervals alone leads to local thermal density matrix at very early times !**
- Hawking-Unruh temperature in Rindler wedge $T(x) = \hbar c/(2\pi x)$

Physics picture

- coherent state vacuum at early time contains entangled pairs of quasi-particles with opposite wave numbers
- on finite rapidity interval $(-\Delta\eta/2, \Delta\eta/2)$ in- and out-flux of quasi-particles with thermal distribution via boundaries
- technically limits $\Delta\eta \rightarrow \infty$ and $M\tau \rightarrow 0$ do not commute
 - $\Delta\eta \rightarrow \infty$ for any finite $M\tau$ gives pure state
 - $M\tau \rightarrow 0$ for any finite $\Delta\eta$ gives thermal state with $T = 1/(2\pi\tau)$

Bosonized massive Schwinger model

[ongoing work with Lara Kuhn, Jürgen Berges]



- scalar theory with potential





$$V(\Phi) = \frac{1}{2}M^2\Phi^2 + J \cos(2\sqrt{\pi}\phi + \theta)$$

- dimensionless coupling strength $g = 2\sqrt{\pi}J/M^2$
- initial value

$$\Phi(0) = \Phi_{\text{vac}} + \sqrt{\pi}$$


Time evolution of background field

- solve equation of motion for background field in expanding geometry
- non-linear oscillations damped by expansion





Excitations

- consider now perturbations around expanding background fields
- linear problem: quantization in time dependent situation
- (quasi-) particle production
- similar: cosmology, strong electric fields
- calculate this for different parameters as function of wave number





Asymptotic particle spectrum

- particles per unit rapidity
- characteristic peaks due to resonance-like phenomena
- for large coupling $g = 2\sqrt{\pi}J/M^2$ exponential decay



Quantum simulation

[ongoing work with Lara Kuhn]



- two one-dimensional Bose-Einstein condensates with tunnel coupling allow to realize model Lagrangian for relative phase $\phi = \varphi_1 - \varphi_2$

$$\mathcal{L} = \frac{1}{2}\dot{\phi}^2 - \frac{c_s^2}{2}(\nabla\phi)^2 - V_0 \left[-\cos(\phi) + \frac{1}{2}\lambda^2 \sin^2(\phi) \right]$$


with velocity of sound c_s

- λ from periodic in time modulation of tunnel coupling [Fialko, Opanchuk, Sidorov, Drummond & Brand (2014)] (challenging to implement)

Tunneling and the bounce

[ongoing work with Lara Kuhn]

- consider field in metastable vacuum
- tunneling can be described by semi-classical methods
- false vacuum decay by bounce solution [S. Coleman (1977)]
- could be induced artificially
- interior of forward sound cone resembles closely expanding string solution



Conclusions

- rapidity intervals in an expanding string are entangled
- at very early times theory effectively conformal

$$\frac{1}{\tau} \gg m, q$$

- entanglement entropy extensive in rapidity $\frac{dS}{d\Delta\eta} = \frac{c}{6}$
- determined by conformal charge $c = N_c \times N_f + 2$
- reduced density matrix for conformal field theory is of locally thermal form with temperature

$$T = \frac{\hbar}{2\pi\tau}$$

- expanding QCD string dynamics could be quantum simulated through two frequency sine-Gordon model

$$V(\phi) \sim \left[-\cos(\phi) + \frac{1}{2}\lambda^2 \sin^2(\phi) \right]$$