


Extended symmetries

Stefan Floerchinger (Friedrich-Schiller-Universität Jena)

ERG 2022 Conference, Berlin
July 25, 2022.



Symmetries



- constrain the form of the microscopic action S
- constrain the form of the quantum effective action Γ
- imply conservation laws [Noether (1918)]
- ... and more ...

$$\Gamma[\mathbf{g}\phi] = \Gamma[\phi]$$

Extended symmetries

- Lie group action on fields with generators T_j

$$\chi(x) \rightarrow \chi(x) + id\xi^j T_j\chi(x)$$

- standard symmetries: action is invariant

$$dS[\chi] = \int_x \frac{\delta S[\chi]}{\delta \chi(x)} id\xi^j T_j\chi(x) = 0$$

- extended symmetries: change in action is *linear* in the fields
[Canet, Delamotte & Wschbor, PRE 91, 053004 (2015)]

$$d\Gamma[\phi] = \langle dS[\chi] \rangle \sim \langle \chi \rangle = \phi$$

- constraints $\Gamma[\phi]$ as powerful as a standard symmetry
- works also when $\delta S[\chi]$ is linear in available composite fields
[Floerchinger & Grossi, PRD 105, 085015 (2022)]

Example: time-dependent Galilei transformations

- non-relativistic fluids
- time-dependent Galilei transformation

$$\mathbf{x} \rightarrow \mathbf{x} + \mathbf{v}T(t), \quad \mathbf{p} \rightarrow \mathbf{p} + m\mathbf{v}\frac{d}{dt}T(t)$$

- changes the action by a term linear in fluid velocity field
- allows to derive a powerful set of Ward identities
- has been used in the context of Navier-Stokes fluid turbulence to obtain closed solutions in certain limits
[Canet, Delamotte & Wschbor, PRE 91, 053004 (2015), PRE 93, 063101 (2016), Tarpin, Canet & N. Wschebor, Phys. Fluids 30, 055102 (2018)]
- can be used in cosmological structure formation to close flow equations
[Erschfeld, Floerchinger, PRD 105, 023506 (2022)]
[Talk by Alaric Erschfeld at this conference]

Noether currents from the quantum effective action

[Floerchinger & Grossi, PRD 105, 085015 (2022)]

- symmetries imply conservation laws
- conserved currents are of interest
- fluid dynamics based on them
- renormalized or full currents can be obtained directly from quantum effective action $\Gamma[\phi]$
- need to couple to external gauge fields
- *extended symmetries* \Rightarrow *non-conserved* Noether currents

Extended symmetries with gauge fields

[Floerchinger & Grossi, PRD 105, 085015 (2022)]

- consider transformation of fields

$$\phi(x) \rightarrow \phi(x) + id\xi^j(x) T_j \phi(x)$$

- might be non-Abelian with structure constants

$$[T_k, T_l] = if_{kl}^j T_j$$

- introduce external gauge field and covariant derivative

$$D_\mu \phi(x) = \left(\nabla_\mu - iA_\mu^j(x) T_j \right) \phi(x)$$

- gauge field transforms as usual

$$A_\mu^j(x) \rightarrow A_\mu^j(x) + f_{kl}^j A_\mu^k(x) d\xi^l(x) + \nabla_\mu d\xi^j(x)$$

Extended conservation laws

[Floerchinger & Grossi, PRD 105, 085015 (2022)]

- change of effective action $\Gamma[\phi, A]$

$$\Gamma[\phi + id\xi^j T_j \phi, A_\mu^j + f_{kl}^j A_\mu^k d\xi^l + \nabla_\mu d\xi^j] = \Gamma[\phi] + \int d^d x \sqrt{g} \left\{ \mathcal{I}_j(x) d\xi^j(x) \right\}$$

- define current through

$$\mathcal{J}_j^\mu(x) = \frac{1}{\sqrt{g}} \frac{\delta \Gamma}{\delta A_\mu^j(x)}$$

- obtain conservation-type relation (for $\delta\Gamma/\delta\phi = 0$)

$$D_\mu \mathcal{J}_j^\mu(x) = \nabla_\mu \mathcal{J}_j^\mu(x) + f_{jk}^l A_\mu^k(x) \mathcal{J}_l^\mu(x) = -\mathcal{I}_j(x)$$

- standard symmetry $\mathcal{I}_j(x) = 0 \Rightarrow$ conserved Noether current
- extended symmetry $\mathcal{I}_j(x) \neq 0 \Rightarrow$ non-conserved Noether current
- right hand side is known, therefore still very useful !

Space-time symmetries and their extensions

Why curved space?

- spacetime metric $g_{\mu\nu}(x)$ provides source for $T^{\mu\nu}(x)$
- metric is actually a gauge field
- full renormalized $T^{\mu\nu}(x)$ follows from variation of $\Gamma[\phi, g]$
- can still evaluate everything in flat space in the end

Covariant energy-momentum conservation

- quantum effective action $\Gamma[\phi, g]$ at stationary matter fields

$$\frac{\delta}{\delta\phi(x)}\Gamma[\phi, g] = 0$$

- energy-momentum tensor defined by

$$\delta\Gamma[\phi, g] = \frac{1}{2} \int d^d x \sqrt{g} T^{\mu\nu}(x) \delta g_{\mu\nu}(x)$$

- diffeomorphism is gauge transformation of metric

$$g_{\mu\nu}(x) \rightarrow g_{\mu\nu}(x) + \nabla_\mu \varepsilon_\nu(x) + \nabla_\nu \varepsilon_\mu(x)$$

- from invariance of $\Gamma[\phi, g]$ under diffeomorphisms

$$\nabla_\mu T^{\mu\nu}(x) = 0$$

- work here in Riemann geometry with Levi-Civita connection

$$\delta\Gamma_{\mu}{}^{\rho}{}_{\nu} = \frac{1}{2} g^{\rho\lambda} (\nabla_\mu \delta g_{\nu\lambda} + \nabla_\nu \delta g_{\mu\lambda} - \nabla_\lambda \delta g_{\mu\nu})$$

Relativistic fluid dynamics

Energy-momentum tensor and conserved current

$$T^{\mu\nu} = \epsilon u^\mu u^\nu + (p + \pi_{\text{bulk}})\Delta^{\mu\nu} + \pi^{\mu\nu}$$

$$N^\mu = n u^\mu + \nu^\mu$$

- tensor decomposition using fluid velocity u^μ , $\Delta^{\mu\nu} = g^{\mu\nu} + u^\mu u^\nu$
- thermodynamic equation of state $p = p(T, \mu)$

Covariant **conservation laws** $\nabla_\mu T^{\mu\nu} = 0$ and $\nabla_\mu N^\mu = 0$ imply

- equation for energy density ϵ
- equation for fluid velocity u^μ
- equation for particle number density n

Need **further evolution equations** [e.g Israel & Stewart]

- equation for shear stress $\pi^{\mu\nu}$
- equation for bulk viscous pressure π_{bulk}

$$\tau_{\text{bulk}} u^\mu \partial_\mu \pi_{\text{bulk}} + \dots + \pi_{\text{bulk}} = -\zeta \nabla_\mu u^\mu$$

- equation for diffusion current ν^μ
- non-hydrodynamic degrees of freedom are needed for relativistic causality!

Why non-Riemannian geometry?

- can one learn more by studying further deformations of geometry?
- some equations of motion for non-ideal fluids still missing
- one can still evaluate everything in Riemannian geometry / flat space in the end

Local scaling or Weyl gauge transformations

- transforms matter fields

$$\phi(x) \rightarrow e^{-\Delta\phi\zeta(x)}\phi(x)$$

- scales metric

$$g_{\mu\nu}(x) \rightarrow e^{2\zeta(x)}g_{\mu\nu}(x)$$

- Weyl gauge field (Abelian)

$$B_\mu(x) \rightarrow B_\mu(x) - \partial_\mu\zeta(x)$$

- variation of effective action with respect to $B_\mu(x)$ gives a current $W^\mu(x)$
- equation of motion from variation with respect to $\zeta(x)$

$$\nabla_\rho W^\rho(x) = \frac{2}{d} [T^\mu{}_\mu(x) - \mathcal{U}^\mu{}_\mu(x)]$$

- in general not conserved but right hand side can be calculated
- vanishes for conformal field theories in flat space

Non-Riemannian geometry

[Floerchinger & Grossi, PRD 105, 085015 (2022)]

- general connection

$$\Gamma_{\mu\sigma}^{\rho} = \frac{1}{2}g^{\rho\lambda} (\partial_{\mu}g_{\sigma\lambda} + \partial_{\sigma}g_{\mu\lambda} - \partial_{\lambda}g_{\mu\sigma}) + C_{\mu\sigma}^{\rho} \\ + \hat{B}_{\mu\sigma}^{\rho} + \hat{B}_{\sigma\mu}^{\rho} - \hat{B}_{\mu\sigma}^{\rho} + B_{\mu}\delta^{\rho}_{\sigma} + B_{\sigma}\delta_{\mu}^{\rho} - B^{\rho}g_{\mu\sigma}$$

- **contorsion** $C_{\mu\sigma}^{\rho}$ = gauge field for local Lorentz transformations
- **Weyl gauge field** B_{μ} = gauge field for local dilatations
- **proper non-metricity** $\hat{B}_{\mu\sigma}^{\rho}$ = gauge field for local shear transformations
- together form the group $GL(d)$ of basis changes in tangent space / the frame bundle

Hypermomentum current

[von der Heyde, Kerlick & Hehl (1976)]

[Floerchinger & Grossi, PRD 105, 085015 (2022)]

- in Non-Riemannian geometry, affine connection $\Gamma_{\mu}^{\rho}{}_{\sigma}(x)$ can be varied independent of the metric $g_{\mu\nu}(x)$

$$\delta\Gamma = \int d^d x \sqrt{g} \left\{ \frac{1}{2} \mathcal{U}^{\mu\nu}(x) \delta g_{\mu\nu}(x) - \frac{1}{2} \mathcal{S}^{\mu}{}_{\rho}{}^{\sigma}(x) \delta \Gamma_{\mu}^{\rho}{}_{\sigma}(x) \right\}$$

with new symmetric tensor $\mathcal{U}^{\mu\nu}$ and *hypermomentum* current $\mathcal{S}^{\mu}{}_{\rho}{}^{\sigma}$

- hypermomentum current can be decomposed further

$$\mathcal{S}^{\mu}{}_{\rho}{}^{\sigma} = Q^{\mu}{}_{\rho}{}^{\sigma} + W^{\mu} \delta_{\rho}^{\sigma} + S^{\mu}{}_{\rho}{}^{\sigma} + S^{\sigma\mu}{}_{\rho} + S_{\rho}{}^{\mu\sigma}$$

with

- spin current $S^{\mu\rho\sigma} = -S^{\mu\sigma\rho}$
- dilatation current W^{μ}
- shear current $Q^{\mu\rho\sigma} = Q^{\mu\sigma\rho}, \quad Q^{\mu\rho}{}_{\rho} = 0$

Equations of motion for shear current

[Floerchinger & Grossi, PRD 105, 085015 (2022)]

- variation of connection contains Levi-Civita part and non-Riemannian part

$$\delta\Gamma_{\mu}{}^{\rho}{}_{\sigma} = \frac{1}{2}g^{\rho\lambda}(\nabla_{\mu}\delta g_{\sigma\lambda} + \nabla_{\sigma}\delta g_{\mu\lambda} - \nabla_{\lambda}\delta g_{\mu\sigma}) + \delta C_{\mu}{}^{\rho}{}_{\sigma} + \delta D_{\mu}{}^{\rho}{}_{\sigma}$$

- variation at $\delta C_{\mu}{}^{\rho}{}_{\sigma} = \delta D_{\mu}{}^{\rho}{}_{\sigma} = 0$ gives energy-momentum tensor

$$T^{\mu\nu} = \mathcal{U}^{\mu\nu} + \frac{1}{2}\nabla_{\rho}(Q^{\rho\mu\nu} + W^{\rho}g^{\mu\nu})$$

- new equation of motion for shear current

$$\nabla_{\rho}Q^{\rho\mu\nu} = 2\left[T^{\mu\nu} - \mathcal{U}^{\mu\nu} - \frac{g^{\mu\nu}}{d}(T^{\sigma}{}_{\sigma} - \mathcal{U}^{\sigma}{}_{\sigma})\right]$$

Spin current

- tetrad formalism: vary tetrad V_μ^A and spin connection Ω_μ^{AB}

$$\delta\Gamma = \int d^d x \sqrt{g} \left\{ \mathcal{T}^\mu_A(x) \delta V_\mu^A(x) - \frac{1}{2} S^\mu_{AB}(x) \delta \Omega_\mu^{AB}(x) \right\}$$

with

- canonical energy-momentum tensor \mathcal{T}^μ_A
- spin current S^μ_{AB}
- symmetric energy-momentum tensor in Belinfante-Rosenfeld form

$$T^{\mu\nu}(x) = \mathcal{T}^{\mu\nu}(x) + \frac{1}{2} \nabla_\rho [S^{\rho\mu\nu}(x) + S^{\mu\nu\rho}(x) + S^{\nu\mu\rho}(x)]$$

- equation of motion for spin current

$$\nabla_\mu S^{\mu\rho\sigma} = \mathcal{T}^{\sigma\rho} - \mathcal{T}^{\rho\sigma}$$

- non-conserved Noether current

Example: Scalar field

- action for scalar field in d spacetime dimensions

$$\Gamma = \int d^d x \sqrt{g} \left\{ -\frac{1}{2} g^{\mu\nu} \bar{\nabla}_\mu \varphi \bar{\nabla}_\nu \varphi - U(\varphi) - \frac{1}{2} \xi R \varphi^2 \right\},$$

- co-covariant derivative

$$\bar{\nabla}_\mu \varphi = (\partial_\mu - \Delta_\varphi B_\mu) \varphi = \left(\partial_\mu - \frac{d-2}{2} B_\mu \right) \varphi$$

- how does this extend to non-Riemannian space? Is R replaced by the generalized Riemann scalar \bar{R} ?
- in that case one finds hypermomentum current

$$\mathcal{S}^\mu{}_\rho{}^\sigma = -\frac{d-2}{2d} \delta^\sigma{}_\rho \partial^\mu \varphi^2 - \xi g^{\mu\sigma} \partial_\rho \varphi^2 + \xi \delta^\mu{}_\rho \partial^\sigma \varphi^2.$$

- Weyl current (vanishes for conformal choice of ξ)

$$W^\mu = \left(\xi \frac{2d-2}{d} - \frac{d-2}{2d} \right) \partial^\mu \varphi^2$$

Implications for relativistic fluid dynamics

- dilatation current, shear current and spin current provide additional information about quantum fields out-of-equilibrium
- their contribution to energy-momentum tensor comes with derivatives and vanishes in equilibrium or for ideal fluids
- new equations of motion for “non-hydrodynamic” modes
- can be useful to set relativistic fluid dynamic on more solid basis

Conclusions

- extended notion of symmetries
- more Ward identities that sometimes allow to close functional renormalization group equations
- extended notion of Noether currents
- more conservation-type equations of motion that are useful for non-equilibrium dynamics
- studying quantum field theory in non-Riemannian geometry can be useful
- dilatation current, shear current and spin current
- new equations of motion for fluids
- to do: extension to fluids with additional charges